

Technical Procedures Bulletin

**Subject: Wave Forecasting
for Eastern North Pacific
and Adjacent Waters**

Series No. 491

June 6, 2003

Science Division, Silver Spring, MD 20910

FIRST TPB ABOUT THIS PRODUCT

This bulletin, prepared by Y. Y. Chao, L. D. Burroughs, and H. L. Tolman of the Marine Modeling and Analysis Branch (MMAB), Environmental Modeling Center (EMC), National Centers for Environmental Prediction (NCEP), describes automated wave guidance for Eastern North Pacific and adjacent waters in alphanumeric, and Gridded Binary (GRIB) formats. This guidance was implemented operationally in May 2002.

The Eastern North Pacific and Adjacent Waters regional wave model (ENP) was developed to provide wind wave forecast guidance for the U.S. West coast and Hawaiian waters. The ENP is based on the NOAA WAVEWATCH-III (NWW3) which is described in detail in Technical Procedures Bulletin (TPB) 494 (Chen, Burroughs, and Tolman 2003) and Tolman (2002). The NWW3 provides the boundary conditions to the ENP. The domain extends from 170.25°W to 76.75°W and from 4.75°N to 62.25°N with a grid resolution of 0.25° by 0.25° in latitudinal and longitudinal directions.

Various graphics and text products for the ENP are available at <http://polar.wwb.noaa.gov/waves>, and available for anonymous ftp at <ftp://polar.wwb.noaa.gov/pub/waves>.

The following wind and wave parameters are available in GRIB format at the web site above and will be on AWIPS as GRIB bulletins when AWIPS Build OB2 is distributed to the field: H_s , D_m , T_m , peak wave period and direction, wind sea peak period and direction, wind speed and direction, and u- and v-wind components.

Spectral text bulletins for the ENP are available at the web site above. These files are in ASCII and are available by anonymous ftp from the directory given above. These bulletins became available on the Satellite Broadcast Network (SBN) in May 2003.

The ENP wave guidance is generated four times daily from the 0000, 0600, 1200 and 1800 UTC cycles of the Global Forecast System (GFS).



LeRoy Spayd
Chief, Training and Professional
Development Core



WAVE FORECASTING FOR THE EASTERN NORTH PACIFIC and ADJACENT WATERS

By

Y.Y. Chao, L.D. Burroughs, and H.L. Tolman⁽¹⁾⁽²⁾

1. Introduction

In order to predict wave conditions adequately over the continental shelf and near land boundaries, a regional model which has higher resolution in grid space and possibly in spectral components is required. The regional model also must calculate rigorously the effects of submarine bottom conditions and any currents which may exist on wave growth, transformation and dissipation. A global-scale wave model usually is designed only to provide the general wave pattern over the deep ocean. It does not provide information accurate enough to describe small-scale, complex wave patterns near the coastal areas.

The Eastern North Pacific (ENP) regional wave model is designed to fill the needs of the Western and Pacific Regions, Tropical Prediction Center, and the Ocean Prediction Center (OPC). The model structure is essentially the same as the Western North Atlantic (WNA) regional model which is described in detail in Technical Procedures Bulletin (TPB) 495 (Chao, Burroughs and Tolman, 2003). The NWW3 provides the boundary conditions to the ENP. The domain extends from 170.25°W to 76.75°W and from 4.75°N to 62.25°N with a grid resolution of 0.25° by 0.25° in latitudinal and longitudinal directions.

Various graphics and text products for the ENP are available at

<http://polar.wwb.noaa.gov/waves>,

and available for anonymous ftp at

<ftp://polar.wwb.noaa.gov/pub/waves> .

The following wind and wave parameters are available in GRIB format at the web site above and will be on AWIPS as GRIB bulletins, when software build OB2 is distributed to the field: H_s , D_m , T_m , peak wave period and direction, wind sea peak period and direction, wind speed and direction, and u- and v-wind components.

Spectral text bulletins for the ENP are available at the web site above. These files are in ASCII and are available by anonymous ftp from the ftp site above. These bulletins will be available on the SBN in May 2003, but with a condensed format necessitated by the capabilities of the communications gateway and display capabilities of AWIPS. See fig. 1 for a sample bulletin and Table 1 for the list of points having spectral wave bulletins, their locations, and their bulletin

headers. The ENP wave guidance is generated four times daily out to 168 hours based on the 0000, 0600, 1200 and 1800 UTC cycles of the of the Global Forecast System (GFS).

The ENP model has been run continuously on the IBM computers since September 2001. In the succeeding sections, the established forecasting system along with the model structure will be described, followed by a comparison of predicted results with buoy data and NWW3 model output for identical locations. Finally, the strength of the new model (ENP) and the available products and dissemination is presented.

2. Model Description

Regional wave forecasts for Eastern North Pacific and Hawaiian waters are generated at NCEP by using the ENP model. Fields of directional frequency spectra in 24 directions and 25 frequencies are generated at one hour intervals up to 168 hours. The 24 directions begin at 90 degrees to the east and have a directional resolution of 15 degrees. The 25 frequencies used by the ENP are given by bin in Table 2.

Figure 2 shows the domain of interest and the depth field which is derived from bathymetric data available from the National Geophysical Data Center. Required input wave spectral data for the boundary grid points of the ENP are obtained by linearly interpolating the output of neighboring grids of the NWW3. The wind fields driving the model are obtained from the output of NCEP's operational GFS (Kanamitsu *et al.* 1991 and Caplan *et al.* 1997).

The wind fields are constructed directly from spectral coefficients of the lowest sigma level at $0.5^\circ \times 0.5^\circ$ longitude and latitude resolution, and are interpolated to the resolution of the wave model grid. They are converted to 10 m winds by using a logarithmic profile corrected for stability with air-sea temperature differences. Air and sea temperature data are obtained from the lowest sigma level air temperatures of the GFS and from the SST analysis available in the model.

Finally, the wave model incorporates a dynamically updated ice coverage field in the region. These data are obtained from NCEP's operational automated passive microwave sea ice concentration analysis (Grumbine 1996; updated daily). Ocean currents are not considered in the model at the present.

The model runs four times daily for the 0000, 0600, 1200 and 1800 UTC cycles. Global Data Assimilation System (GDAS) wind fields from the previous 12 hours at 3-h intervals (analyses and 3-h forecasts) are used for a 12-h wave hindcast. Winds from the AVN at 3-h intervals out to 168 hours are used to produce wave forecasts up to 168 hours at hourly intervals.

3. Performance Evaluation

The ENP has run on the IBM mainframe computer continuously since September 2001. The adequacy of the model has been evaluated by comparing model output of the ENP with observed data at NDBC buoys. Figure 3 shows the locations of buoy measurements used for verifying model output in the vicinity of Hawaiian Islands. The model data at the buoy station are obtained based on interpolating gridded data surrounding the buoy station.

Figure 4 shows the scatter plot of the hindcast significant wave height NWW3 and ENP for December 2001 through January 2002 . Also shown are the following statistical indices: the mean bias error (bias), standard deviation(std),root mean square error (rms), scatter index (SI), the slope of best fit line and correlation coefficient (c.c). The total number of data points (4116) used in the analysis combines the results from the measurement of four buoys located in deep water. It can be seen from the figure that the ENP provides comparable or slightly better statistical results than NWW3.

An important aspect of the island effect on waves, which can only be detected by using a high grid resolution model, is shown in Fig. 5. In Fig. 5 , the wave height time series of measurements (+ marks), NWW3 (red lines) and ENP (green lines) at buoy locations 51001 and 51004 are presented. During this period of time, the dominant wave direction was from north-west to south-east, closely parallel to the alignment of the group of Hawaiian islands . At Buoys 51001, which is located north-westward from the islands, the wave height of NWW3 and ENP are essentially the same. At buoy 51004 which was located south-eastward from the islands, however, over-predictions of the wave height by the NWW3 model are clearly indicated. In contrast, the ENP model predictions are very close to buoy measurements. The differences demonstrated here have been greatly reduced by the introduction of sub-grid islands in the wave model suite.

Figure 6 shows available buoy measurements off and along the coastal line of West Coast. The NWW3 and ENP model predictions at those deep water buoys are virtually the same. The statistical evaluation of ENP hindcast against the buoy measurements in the shelf is shown in Fig. 7. The performance evaluation of NWW3 model against these shelf buoys cannot be made because of coarse grid mesh so that an adequate interpolation is not possible. The overall performance of the ENP model over the shelf during the five month period (September 2001 through January 2001) is very satisfying.

4. Available Products and Dissemination

The following wind and wave parameters are available in GRIB format at

<http://polar.wwb.noaa.gov/waves>

and on AWIPS as GRIB bulletins: H_s , D_m , T_m , peak wave period and direction, wind sea peak period and direction, wind speed and direction, and u- and v-wind components. Spectral text bulletins are also available on the web at the site above and will be on AWIPS as soon AWIPS software build OB2 is distributed to the field.

a. GRIB bulletins

GRIB bulletins will be available for use in AWIPS when software build OB2 is distributed to the field later this year. Table 3 gives the bulletin headers and their meaning. Bulletins are available at 6-h intervals from 00- through 72-h and at 12-h intervals from 72- through 168-h. Available parameters are H_s , D_m , T_m , peak wave period and direction, wind sea peak wave period and direction, and u and v components of the wind velocity. A 0.25 x 0.25 degree lon./lat. grid is used with a domain of 170.25°W - 76.75°W by 4.75°N - 62.25°N.

b. Web Based Alphanumeric Spectral Messages

Spectral text bulletins are presented for numerous points of the ENP. These bulletins are in ASCII and are available on the INTERNET, and became available to the field on the SBN in May 2003. The line length of the web table is 130 characters by 100 lines. The header of the table identifies the output location, the generating model and the run date and cycle of the data presented. At the bottom of the table, a legend is printed. The table consists of 8 columns. The first column gives the time of the model results with a day and hour (the corresponding month and year can be deduced from the header information). The second column presents the overall significant wave height (H_s), the number of individual wave fields with a wave height over 0.15 m that could not be tracked in the table (x). Individual wave fields in the spectrum are identified by using a partitioning scheme similar to that of Gerling (1992). In the remaining six columns individual wave fields identified with their wave height (H_s), peak wave period (T_p) and mean wave direction (dir, direction in which waves travel relative to North). Generally, each separate wave field is tracked in its own column. Such tracking, however is not guaranteed to work all the time. An asterisk (*) in a column identifies that the wave field is at least partially under the influence of the local wind, and, therefore, most likely part of the local wind sea. All other wave fields are pure swell.

b. Spectral text bulletins for AWIPS

The format for the spectral text bulletins sent to AWIPS is generally the same as that for the web, except that the period is to the nearest second, the wave heights are to the nearest foot, the direction is from (meteorological, rather than oceanographic), the number of fields that couldn't be tracked is not given, and the asterisk indicating when a wave field is, at least, partially under the influence of the local wind is not shown. The bulletin width is 69 characters, which is a legacy of the teletype era and the display capability of AWIPS. A sample bulletin is shown in fig. 1, and the list of points for the NWW3 is given in Table 1.

5. References

- Booij, N. and L.H. Holthuijsen, 1987: Propagation of ocean waves in discrete spectral wave models. *J. Comput. Phys.*, **68**, 307-326.
- Caplan, P., J. Derber, W. Gemmill, S.-Y. Hong, H.-L. Pan and D. Parish, 1997: Changes to the NCEP operational medium-range forecast model analysis/forecast system. *Wea. Forecasting*, **12**, 581-594.
- Chalikov, D.V. and Belevich, M.Y., 1993: One-dimensional theory of the boundary layer. *Boundary-Layer Meteor.*, **63**, 65-96.
- Chao, Y.Y., L.D. Burroughs and H.L. Tolman, 2003: Wave Forecasting for Western North Atlantic Ocean and Adjacent Waters. *Technical Procedures Bulletin No. 495*, National Weather Service, NOAA, U.S. Department of Commerce, [Available At <http://polar.wwb.noaa.gov/mmab/tpbs/wnatpb> .]
- Chen, H.S., L.D. Burroughs and H.L. Tolman, 2003: Ocean Surface Waves Model WAVEWATCH III. *Technical Procedures Bulletin No. 494*, National Weather Service, NOAA, U.S. Department of Commerce. [Available at <http://polar.wwb.noaa.gov/mmab/tpbs/tpb494>.]

Gerling, T.W., 1992: Partitioning sequences and arrays of directional wave spectra into component systems. *J. Atmos. Ocean. Technol.*, **9**, 444-458.

Grumbine, R.W., 1996: Automated passive microwave sea ice concentration analysis at NCEP. *Ocean Modeling Branch Tech Note No. 120*, NCEP, National Weather Service, NOAA, U.S. Department of Commerce, 13 pp.

Kanamitsu, M., J.C. Alpert, K.A. Campana, P.M. Caplan, D.G. Deaven, M. Iredell, B. Katz, H.-L. Pan, J.E. Sela and G. H. White, 1991: Recent Changes implemented into the global forecast system at NMC. *Wea. Forecasting*, **6**, 425-435.

The SWAMP Group, 1985: *Ocean Wave Modeling*. Plenum Press, New York, 256 pp.

Tolman, H.L. and D. Chalikov, 1996: Source terms in a third-generation wind-wave model. *J. Phys. Oceanogr.*, **26**, 2497-2518.

Tolman, H.L., 2002: User manual and system documentation of WAVEWATCH-III version 2.22. *Technical Note No. 222*, Marine Modeling and Analysis Branch, NCEP, National Weather Service, NOAA, U.S. Department of Commerce, 139 pp. [Available at http://polar.wwb.noaa.gov/mmab/papers/tn222/MMAB_222.pdf].

1. H. L. Tolman is a contractor with SAIC.

2. OMB Contribution No. ???

Table 1. Station name, position, and bulletin header for spectral text bulletins at points from the ENP regional wave model.

Station Name	Position (N and W, except where indicated)		AWIPS and WMO Header
	Latitude	Longitude	
Points for Wave Spectra from the Eastern North Pacific (ENP) Regional Wave Model			
Eastern Pacific Points			
46002	42.50	130.30	AGPZ46 KWBJ OSBE01
46006	40.90	137.50	AGPZ46 KWBJ OSBE02
46059	38.00	130.00	AGPZ46 KWBJ OSBE03
46011	34.88	120.87	AGPZ46 KWBJ OSBE04
46012	34.45	122.70	AGPZ46 KWBJ OSBE05
46013	38.23	123.33	AGPZ46 KWBJ OSBE06
46014	39.22	123.97	AGPZ46 KWBJ OSBE07
46022	40.72	124.52	AGPZ46 KWBJ OSBE08
46023	34.71	120.97	AGPZ46 KWBJ OSBE09
46026	37.75	122.82	AGPZ46 KWBJ OSBE10
46027	41.85	124.38	AGPZ46 KWBJ OSBE11
46028	35.74	121.89	AGPZ46 KWBJ OSBE12
46030	40.50	124.50	AGPZ46 KWBJ OSBE13
46042	36.75	122.42	AGPZ46 KWBJ OSBE14
46047	32.43	119.53	AGPZ46 KWBJ OSBE15
46050	44.62	124.53	AGPZ46 KWBJ OSBE16
46062	35.10	121.01	AGPZ46 KWBJ OSBE17

46063	34.25	120.66	AGPZ46 KWB OSBE18
TPC50	30.00	118.00	AGPZ46 KWB OSBE19
TPC51	20.00	135.00	AGPZ46 KWB OSBE20
TPC52	20.00	117.00	AGPZ46 KWB OSBE21
TPC53	06.00	120.00	AGPZ46 KWB OSBE22
TPC54	15.00	95.00	AGPZ46 KWB OSBE23
TPC55	09.00	88.00	AGPZ46 KWB OSBE24
TPC56	06.80	80.00	AGPZ46 KWB OSBE25
46025	33.75	119.08	AGPZ46 KWB OSBE26
46053	34.24	119.85	AGPZ46 KWB OSBE27
46054	34.27	120.45	AGPZ46 KWB OSBE28
SGX01	32.64	117.75	AGPZ46 KWB OSBE29
Eastern Gulf of Alaska Points			
46004	50.97	135.80	AGGA47 KWB OSBE01
46184	53.90	138.87	AGGA47 KWB OSBE02
46205	51.17	134.33	AGGA47 KWB OSBE03
46207	50.86	129.91	AGGA47 KWB OSBE04
46208	52.50	132.70	AGGA47 KWB OSBE05
46083	58.25	138.00	AGGA47 KWB OSBE06
46084	56.39	136.16	AGGA47 KWB OSBE07
46082	59.61	143.67	AGGA47 KWB OSBE08

Canadian Points			
46005	46.10	131.00	AGPZ47 KWB OSBE01
46029	46.12	124.50	AGPZ47 KWB OSBE02
46036	48.35	133.92	AGPZ47 KWB OSBE03
46041	47.34	124.67	AGPZ47 KWB OSBE04
46132	49.73	127.92	AGPZ47 KWB OSBE05
46206	48.84	126.00	AGPZ47 KWB OSBE06
Western Gulf of Alaska Points			
46001	56.30N	148.30W	AGGA48 KWB OSBE01
46066	52.65	155.00	AGGE48 KWB OSBE02
46080	58.00	150.00	AGGE48 KWB OSBE03
Hawaiian Points			
51001	23.40	162.30	AGHW40 KWB OSBE01
51002	17.20	157.80	AGHW40 KWB OSBE02
51003	19.10	160.80	AGHW40 KWB OSBE03
51004	17.40	152.50	AGHW40 KWB OSBE04
HNL01	24.00	158.00	AGHW40 KWB OSBE05
HNL02	22.50	153.00	AGHW40 KWB OSBE06
HNL10	22.00	157.75	AGHW40 KWB OSBE07
HNL11	21.00	158.25	AGHW40 KWB OSBE08
HNL12	19.75	156.50	AGHW40 KWB OSBE09

Notes:

The WMO/AWIPS headers follow the form given for oceanographic data, i.e., $AGA_1A_2i_1i_2$, where i_1 is 4 and always means spectral wave data.

i_2 is the geographic location, where:

- 0 means Pacific Ocean, particularly in proximity to U.S. held islands (Hawaii and Guam's areas of responsibility)
- 1 means proximity to NE Atlantic States from Virginia northward
- 2 means proximity to SE Atlantic States from North Carolina southward and Puerto Rico
- 4 means proximity to southern Gulf of Mexico states
- 6 means proximity to Pacific States and southern British Columbia
- 7 means proximity to Panhandle of Alaska and northern British Columbia (Juneau's areas of responsibility)
- 8 means proximity to southern and southwestern Alaska (Anchorage's areas of responsibility)

A_1A_2 is used by the originating office (NCEP/NCO) to identify the oceanic area of the point, where:

- NT - Western Atlantic
- GX - Gulf of Mexico
- CA - Caribbean Sea
- PZ - Eastern Pacific
- GA - Gulf of Alaska
- PN - North Pacific including Bering Sea
- AC - Arctic Ocean
- HW - Hawaiian Waters
- PW - Western Pacific
- XT - Tropical Belt
- PS - South Pacific

The AWIPS identifier form is $NNNxxx$: where NNN is OSB - Oceanographic Spectral Bulletin, and xxx takes the form: mnn - where m is the wave model and nn is the number of the point in a given geographic location according to note 2 above. nn can range from 01 - 99.

m is the wave model where: N is the NOAA WAVEWATCH III global wave model

- A is the Alaska Waters regional wave model
- W is the Western North Atlantic regional wave model
- H is the North Atlantic Hurricane regional wave model
- E is the Eastern North Pacific regional wave model
- P is the Eastern Pacific Hurricane regional wave model
- X is the Western North Pacific regional wave model
- T is the Western Pacific Typhoon regional wave model

Table 2. The center frequencies and corresponding band widths with center period by frequency bin.

bin number	center frequency (Hz)	frequency band width (Hz)	center period (s)
1	.0418	.00399	23.94
2	.0459	.00439	21.76
3	.0505	.00482	19.79
4	.0556	.00531	17.99
5	.0612	.00584	16.35
6	.0673	.00642	14.87
7	.0740	.00706	13.51
8	.0814	.00777	12.29
9	.0895	.00855	11.17
10	.0985	.00940	10.15
11	.1083	.01034	9.23
12	.1192	.01138	8.39
13	.1311	.01251	7.63
14	.1442	.01376	6.93
15	.1586	.01514	6.30
16	.1745	.01666	5.73
17	.1919	.01832	5.21
18	.2111	.02015	4.74
19	.2322	.02217	4.31
20	.2555	.02438	3.91
21	.2810	.02682	3.56
22	.3091	.02951	3.24
23	.3400	.03246	2.94
24	.3740	.03570	2.67
25	.4114	.03927	2.43

Table 3. WMO GRIB bulletin header descriptors.

T_1	T_2^1	A_1^2	A_2	dd	Station id
O	A B C J K M N P Y	M	A C E G I J K L M X N Y O P Q Z R S T U	88	KWBJ

Where:

- T_1 is the bulletin type descriptor: O - oceanographic.
- T_2 is the parameter descriptor, see notes below.
- A_1 is the grid and domain descriptor: M - 0.25° x 0.25° lon/lat grid over domain from 98.25°W to 29.75°W and from 0.25°S to 50.25°N..
- A_2 is the forecast hour descriptor, see notes below.
- dd is the surface descriptor: 88 - ocean surface.

Notes:

1. Parameter descriptors

- A - u-wind component
- B - v-wind component
- C - Total significant wave height
- J - Peak wave period
- K - Peak wave direction
- M - Peak wind sea period
- N - peak wind sea direction
- P - $D_m Y - T_m$

2. Forecast hour descriptors at 6-h intervals from 0- to 72-h and at 12-h intervals from 72- to 168-h.

AGGA47 KWBK 080421
 OSBE01
 LOCATION : 46004 (50.97N 135.80W)
 MODEL : EASTERN NORTH PACIFIC 0.25 DEG
 CYCLE : 20020908 T00Z

<
 DDHH HS SS PP DDD SS PP DDD SS PP DDD SS PP DDD SS PP DDD SS PP DDD

```

-----
0712 5 3 06 356 2 10 276 2 15 197 1 12 204 1 08 219 1 20 201
0713 5 3 05 356 2 10 276 2 15 197 1 12 204 1 08 219 1 19 201
0714 5 2 10 276 2 15 197 1 12 204 1 08 219 1 19 201
0715 4 2 06 340 2 10 276 2 15 197 1 12 204 1 08 219 1 19 201
0716 4 2 06 340 2 10 276 2 15 197 1 12 204 1 08 219 1 19 202
0717 4 2 06 339 2 10 276 2 15 199 1 12 204 1 08 220 1 18 198
0718 4 2 05 340 2 10 276 2 15 199 1 12 204 1 08 220 1 18 198
0719 4 2 05 341 2 10 277 2 15 199 1 12 204 1 08 220 1 18 199
0720 4 2 05 341 2 10 277 2 15 199 1 12 204 1 08 220 1 18 199
0721 4 2 05 341 2 10 277 2 15 199 1 12 204 1 08 220 2 18 200
0722 4 2 05 340 2 10 277 2 15 199 1 12 204 1 08 220 2 18 200
0723 4 2 05 338 2 10 277 2 15 199 1 12 204 1 08 220 2 18 200
0800 4 2 05 337 2 10 277 2 15 199 1 12 204 1 08 219 2 18 201
0801 4 2 05 336 2 09 277 2 15 199 1 12 203 1 08 219 2 18 201
0802 4 2 05 334 2 09 277 2 15 199 1 12 203 1 08 219 2 18 201
0803 4 2 09 278 2 15 201 1 07 219 2 18 201
0804 4 2 05 002 2 09 278 2 15 201 1 03 300 1 07 219 2 18 202
0805 4 2 05 004 2 09 278 2 15 201 1 03 301 1 07 219 2 18 202
0806 4 2 05 005 2 09 278 2 15 201 1 03 299 1 07 219 2 18 202
0807 4 2 05 008 2 09 278 2 15 201 2 03 296 1 07 219 2 18 202
0808 4 2 05 009 2 09 278 2 15 201 2 03 294 1 07 220 2 18 202
0809 4 2 05 009 2 09 277 2 15 201 2 03 289 1 07 219 2 18 202
0810 4 1 06 010 2 09 277 2 15 201 2 04 287 1 07 219 2 18 202
0811 5 1 06 010 2 09 277 2 15 201 2 04 285 1 07 219 2 18 203
0812 5 1 06 010 2 09 277 2 15 201 3 04 283 1 07 219 2 18 203
-----

```

```

-----
1414 4 1 16 215 1 20 203 3 09 284 2 06 224 2 13 210
1415 4 1 16 215 1 20 203 3 09 284 2 06 223 2 13 210
1416 4 2 16 214 1 19 203 3 09 284 2 06 224 2 13 210
1417 4 2 19 209 3 09 284 2 06 224 2 13 210
1418 4 1 18 164 2 18 208 3 09 284 2 06 224 2 13 210
1419 4 1 18 164 2 18 208 3 09 284 2 06 224 2 13 210
1420 4 1 18 164 2 18 208 2 09 284 2 06 224 2 13 211
1421 4 1 18 164 2 18 207 2 09 284 2 06 224 2 13 211
1422 4 1 18 164 2 18 207 2 09 283 2 06 224 2 13 211
1423 4 1 18 164 2 18 207 1 02 051 2 09 283 2 06 225 2 13 211
1500 4 1 18 164 3 18 208 1 02 051 2 09 283 2 06 224
-----

```

DD = DAY OF MONTH
 HH = HOUR OF DAY
 HS = TOTAL SIGNIFICANT WAVE HEIGHT (FEET)
 SS = SIGNIFICANT WAVE HEIGHT OF SEPARATE SYSTEM (FEET)
 PP = PEAK PERIOD OF SEPARATE SYSTEM (WHOLE SECONDS)
 DDD = MEAN DIRECTION OF SEPARATE SYSTEM (DEGREES/"FROM")

Figure 1. Sample wave spectra bulletin from the ENP. These bulletins give a 12-h hindcast as well as a 168-h forecast at 1-h intervals. Only a portion of the bulletin is shown.

ENP model grid and depths





Figure 2. ENP domain and depth field.

ENP Hawaiian Buoy




Figure 3. Locations buoys used to verify ENP model output in the vicinity of the Hawaiian Islands.

Dec. 2001 Jan. 2002




Figure 4. Scatter plots comparing hindcasts of significant wave heights for the NWW3 (left) and the ENP (right) for the months of December 2001 and January 2002.




Figure 5. Comparison of wave heights observed upstream and downstream of the Hawaiian Islands as forecast by the NWW3 and ENP models and measured by buoys 51001 and 51004. These comparisons were made prior to the implementation of the sub-grid islands to all models which has significantly reduced these differences.

ENP West Coast Buoys




Figure 6. Available buoy measurements off and along the West coast of the U.S.

Sept. 2001 Jan. 2002




Figure 7. Statistical evaluation of ENP hindcast against buoy measurements along the shelf of the West coast.